

## Grades Kindergarten – 6

Kindergarten requirements apply to all students  $\geq 5$  years.

|             |   |
|-------------|---|
| DTaP        | <b>5 doses;</b> 4 doses are acceptable if the 4 <sup>th</sup> dose is given on or after the 4 <sup>th</sup> birthday. DT is only acceptable with a letter stating a medical contraindication to DTaP.   |
| Polio       | <b>4 doses;</b> 4 <sup>th</sup> dose must be given on or after the 4 <sup>th</sup> birthday and $\geq 6$ months after the previous dose, or a 5 <sup>th</sup> dose is required. 3 doses are acceptable if the 3 <sup>rd</sup> dose is given on or after the 4 <sup>th</sup> birthday and $\geq 6$ months after the previous dose. |
| Hepatitis B | <b>3 doses;</b> laboratory evidence of immunity acceptable  |
| MMR         | <b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and the 2 <sup>nd</sup> dose must be given $\geq 28$ days after dose 1; laboratory evidence of immunity acceptable  |
| Varicella   | <b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and 2 <sup>nd</sup> dose must be given $\geq 28$ days after dose 1; a reliable history of chickenpox* or laboratory evidence of immunity acceptable   |

## Grades 7 – 12

In ungraded classrooms, grade 7 requirements apply to all students  $\geq 12$  years.

|             |   |
|-------------|---|
| Tdap        | <b>1 dose;</b> and history of DTaP primary series or age appropriate catch-up vaccination. Tdap given at $\geq 7$ years may be counted, but a dose at age 11-12 is recommended if Tdap was given earlier as part of a catch-up schedule. Td should be given if it has been $\geq 10$ years since Tdap.                            |
| Polio       | <b>4 doses;</b> 4 <sup>th</sup> dose must be given on or after the 4 <sup>th</sup> birthday and $\geq 6$ months after the previous dose, or a 5 <sup>th</sup> dose is required. 3 doses are acceptable if the 3 <sup>rd</sup> dose is given on or after the 4 <sup>th</sup> birthday and $\geq 6$ months after the previous dose. |
| Hepatitis B | <b>3 doses;</b> laboratory evidence of immunity acceptable  |
| MMR         | <b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and the 2 <sup>nd</sup> dose must be given $\geq 28$ days after dose 1; laboratory evidence of immunity acceptable  |
| Varicella   | <b>2 doses;</b> first dose must be given on or after the 1 <sup>st</sup> birthday and 2 <sup>nd</sup> dose must be given $\geq 28$ days after dose 1; a reliable history of chickenpox* or laboratory evidence of immunity acceptable   |

## Campers, staff and volunteers 18 years of age and older

|             |  |
|-------------|--|
| MMR         | <b>2 doses,</b> anyone born in or after 1957. 1 dose, anyone born before 1957 outside the U.S. Anyone born in the U.S. before 1957 is considered immune. Laboratory evidence of immunity to measles, mumps and rubella is acceptable   |
| Varicella   | <b>2 doses,</b> anyone born in or after 1980 in the U.S., and anyone born outside the U.S. Anyone born before 1980 in the U.S. is considered immune. A reliable history of chickenpox or laboratory evidence of immunity is acceptable |
| Tdap        | <b>1 dose;</b> Td should be given if it has been $\geq 10$ years since Tdap  |
| Hepatitis B | <b>3 doses for staff whose responsibilities include first aid;</b> laboratory evidence of immunity is acceptable   |

